

CORRECTION

Open Access



# Correction to: multiple imputation for patient reported outcome measures in randomised controlled trials: advantages and disadvantages of imputing at the item, subscale or composite score level

Ines Rombach<sup>1,2\*</sup>, Alastair M Gray<sup>1</sup>, Crispin Jenkinson<sup>3</sup>, David W Murray<sup>2</sup> and Oliver Rivero-Arias<sup>4</sup>

## Correction

Following publication of the original article [1], the authors reported that the following notation wasn't used consistently. The original article has been corrected.

The correct notation is as follows:

Unit-nonresponse

Item-nonresponse

## Author details

<sup>1</sup>Health Economics Research Centre, Nuffield Department of Population Health, University of Oxford, Oxford, UK. <sup>2</sup>Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, Oxford, UK. <sup>3</sup>Health Services Research Unit, Nuffield Department of Population Health, University of Oxford, Oxford, UK. <sup>4</sup>National Perinatal Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford, UK.

Received: 31 August 2018 Accepted: 24 September 2018

Published online: 16 October 2018

## Reference

1. Rombach I, et al. Multiple imputation for patient reported outcome measures in randomised controlled trials: advantages and disadvantages of imputing at the item, subscale or composite score level. *BMC Med Res Methodol.* 2018;18:87. <https://doi.org/10.1186/s12874-018-0542-6>.

\* Correspondence: [ines.rombach@ndorms.ox.ac.uk](mailto:ines.rombach@ndorms.ox.ac.uk)

<sup>1</sup>Health Economics Research Centre, Nuffield Department of Population Health, University of Oxford, Oxford, UK

<sup>2</sup>Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, Oxford, UK

