

RESEARCH

Open Access



Design of a new Z-test for the uncertainty of Covid-19 events under Neutrosophic statistics

Muhammad Aslam^{*} 

Abstract

Background: The existing Z-test for uncertainty events does not give information about the measure of indeterminacy/uncertainty associated with the test.

Methods: This paper introduces the Z-test for uncertainty events under neutrosophic statistics. The test statistic of the existing test is modified under the philosophy of the Neutrosophy. The testing process is introduced and applied to the Covid-19 data.

Results: Based on the information, the proposed test is interpreted as the probability that there is no reduction in uncertainty of Covid-19 is accepted with a probability of 0.95, committing a type-I error is 0.05 with the measure of an indeterminacy 0.10. Based on the analysis, it is concluded that the proposed test is informative than the existing test. The proposed test is also better than the Z-test for uncertainty under fuzzy-logic as the test using fuzz-logic gives the value of the statistic from 2.20 to 2.42 without any information about the measure of indeterminacy. The test under interval statistic only considers the values within the interval rather than the crisp value.

Conclusions: From the Covid-19 data analysis, it is found that the proposed Z-test for uncertainty events under the neutrosophic statistics is efficient than the existing tests under classical statistics, fuzzy approach, and interval statistics in terms of information, flexibility, power of the test, and adequacy.

Keywords: Uncertainty, Classical statistics, Z test, Simulation, Data

Background

The Z-test is playing an important role in analyzing the data. The main aim of the Z-test is to test the mean of the unknown population in decision-making. The Z-test for uncertainty events is applied to test the reduction in the uncertainty of past events. This type of test is applied to test the null hypothesis that there is no reduction in uncertainty against the alternative hypothesis that there is a significant reduction in uncertainty of past events. The Z-test for uncertainty events uses the information of the

past events for testing the reduction of uncertainty [1]. discussed the performance of the statistical test under uncertainty [2]. discussed the design of the Z-test for uncertainty events [3]. worked on the test in the presence of uncertainty [4]. worked on the modification of non-parametric test. The applications of [5], [6], [7] and [8].

[9] mentioned that “statistical data are frequently not precise numbers but more or less non-precise also called fuzzy. Measurements of continuous variables are always fuzzy to a certain degree”. In such cases, the existing Z-tests cannot be applied for the testing of the mean of population or reduction in uncertainty. Therefore, the existing Z-tests are modified under the fuzzy-logic to deal with uncertain, fuzzy, and vague data [10], [11],

*Correspondence: aslam_ravian@hotmail.com
Department of Statistics, Faculty of Science, King Abdulaziz University,
Jeddah 21551, Saudi Arabia



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

[12], [13], [14], [15], [16], [17], [18], [19] worked on the various statistical tests using the fuzzy-logic.

Nowadays, neutrosophic logic attracts researchers due to its many applications in a variety of fields. The neutrosophic logic counters the measure of indeterminacy that is considered by the fuzzy logic, see [20] [21]. proved that neutrosophic logic is efficient than interval-based analysis. More applications of neutrosophic logic can be seen in [22], [23], [24] and [25] [26]. applied the neutrosophic statistics to deal with uncertain data [27]. and [28] presented neutrosophic statistical methods to analyze the data. Some applications of neutrosophic tests can be seen in [29], [30] and [31].

The existing Z-test for uncertainty events under classical statistics does not consider the measure of indeterminacy when testing the reduction in events. By exploring the literature and according to the best of our knowledge, there is no work on Z-test for uncertainty events under neutrosophic statistics. In this paper, the medication of Z-test for uncertainty events under neutrosophic statistics will be introduced. The application of the proposed test will be given using the Covid-19 data. It is expected that the proposed Z-test for uncertainty events under neutrosophic statistics will be more efficient than the existing tests in terms of the power of the test, information, and adequacy.

Methods

The existing Z-test for uncertainty events can be applied only when the probability of events is known. The existing test does not evaluate the effect of the measure of indeterminacy/uncertainty in the reduction of uncertainty of past events. We now introduce the modification of the Z-test for uncertainty events under neutrosophic statistics. With the aim that the proposed test will be more effective than the existing Z-test for uncertainty events under classical statistics. Let $A_N = A_L + A_U I_{A_N}$; $I_{A_N} \in [I_{A_L}, I_{A_U}]$ and $B_N = B_L + B_U I_{B_N}$; $I_{B_N} \in [I_{B_L}, I_{B_U}]$ be two neutrosophic events, where lower values A_L, B_L denote the determinate part of the events, upper values $A_U I_{A_N}, B_U I_{B_N}$ be the indeterminate part, and $I_{A_N} \in [I_{A_L}, I_{A_U}]$, $I_{B_N} \in [I_{B_L}, I_{B_U}]$ be the measure of indeterminacy associated with these events. Note here that the events $A_N \in [A_L, A_U]$ and $B_N \in [B_L, B_U]$ reduces to events under classical statistics (determinate parts) proposed by [2] if $I_{A_L} = I_{B_L} = 0$. Suppose $n_N = n_L + n_U I_N$; $I_N \in [I_L, I_U]$ be a neutrosophic random sample where n_L is the lower (determinate) sample size and $n_U I_N$ be the indeterminate part and $I_N \in [I_L, I_U]$ be the measure of uncertainty in selecting the sample size. The neutrosophic random sample reduces to random sample if no uncertainty is found

in the sample size. The methodology of the proposed Z-test for uncertainty events is explained as follows.

Suppose that the probability that an event $A_N \in [A_L, A_U]$ occurs (probability of truth) is $P(A_N) \in [P(A_L), P(A_U)]$, the probability that an event $A_N \in [A_L, A_U]$ does not occur (probability of false) is $P(A_N^c) \in [P(A_L^c), P(A_U^c)]$, the probability that an event $B_N \in [B_L, B_U]$ occurs (probability of truth) is $P(B_N) \in [P(B_L), P(B_U)]$, the probability that an event $B_N \in [B_L, B_U]$ does not occur (probability of false) is $P(B_N^c) \in [P(B_L^c), P(B_U^c)]$. It is important to note that sequential analysis is done to reduce the uncertainty by using past events information. The purpose of the proposed test is whether the reduction of uncertainty is significant or not. Let $Z_N \in [Z_L, Z_U]$ be neutrosophic test statistic, where Z_L and Z_U are the lower and upper values of statistic, respectively and defined by.

$$Z_N \in \left\{ \frac{P(B_{+kL}|A_L) - P(B_L)}{\sqrt{\frac{P(B_L)[1-P(B_L)][1-P(A_L)]}{(n_L - k_L)P(A_L)}}}, \frac{P(B_{+kU}|A_U) - P(B_U)}{\sqrt{\frac{P(B_U)[1-P(B_U)][1-P(A_U)]}{(n_U - k_U)P(A_U)}}} \right\} \quad (1)$$

Note that $P(B_{+kN}|A_N) = P(B_N|A_N)$ at lag k_N , where $P(B_N|A_N) \in [P(B_L|A_L), P(B_U|A_U)]$ denotes the conditional probability. It means that the probability of event $P(B_N) \in [P(B_L), P(B_U)]$ will be calculated when the event $A_N \in [A_L, A_U]$ has occurred.

The neutrosophic form of the proposed test statistic, say $Z_N \in [Z_L, Z_U]$ is defined by.

$$Z_N = \frac{P(B_{+kL}|A_L) - P(B_L)}{\sqrt{\frac{P(B_L)[1-P(B_L)][1-P(A_L)]}{(n_L - k_L)P(A_L)}}} + \frac{P(B_{+kU}|A_U) - P(B_U)}{\sqrt{\frac{P(B_U)[1-P(B_U)][1-P(A_U)]}{(n_U - k_U)P(A_U)}}} I_{ZN} \in [I_{ZL}, I_{ZU}] \quad (2)$$

The alternative form of Eq. (2) can be written as.

$$Z_N = (1 + I_{ZN}) \frac{P(B_{+kN}|A_N) - P(B_N)}{\sqrt{\frac{P(B_N)[1-P(B_N)][1-P(A_N)]}{(n_N - k_N)P(A_N)}}}; I_{ZN} \in [I_{ZL}, I_{ZU}] \quad (3)$$

The proposed test $Z_N \in [Z_L, Z_U]$ is the extension of several existing tests. The proposed test reduces to the existing Z test under classical statistics when $I_{ZN} = 0$. The proposed test is also an extension of the Z test under fuzzy approach and interval statistics.

The proposed test will be implemented as follows.

Step-1: state the null hypothesis H_0 : there is no reduction in uncertainty vs. the alternative hypothesis H_1 : there is a significant reduction in uncertainty.

Step-2: Calculate the statistic $Z_N \in [Z_L, Z_U]$

Step-3: Specify the level of significance α and select the critical value from [2].

Step-4: Do not accept the null hypothesis if the value of $Z_N \in [Z_L, Z_U]$ is larger than the critical value.

Results

The application of the proposed test is given in the medical field. The decision-makers are interested to test the reduction in uncertainty of Covid-19 when the measure of indeterminacy/uncertainty is $I_{ZN} \in [0, 0.10]$. The decision-makers are interested to test that the reduction in death due to Covid-19 (event A_N) with the increase in Covid-19 vaccines (event B_N). By following [2], the sequence in which both events occur is given as

$$A_N A_N, B_N A_N, B_N A_N, B_N B_N, A_N B_N, A_N B_N$$

where $n_N \in [12, 12]$, $k_N \in [1, 1]$, $P(A_N) = 6/12 = 0.5$ and $P(B_N) = 6/12 = 0.5$.

Note here that event A_N occurs 6 times and that of these 6 times B_N occurs immediately after A_N five times. Given that A_N has occurred, we get

$(B_{+kN} | A_N) = P(B_N | A_N) = 5/6 = 0.83$ at lag 1. The value of $Z_N \in [Z_L, Z_U]$ is calculated as

$$Z_N = (1 + 0.1) \frac{0.83 - 0.50}{\sqrt{\frac{0.50[1 - 0.50][1 - 0.50]}{(12 - 1)0.50}}} = 2.42; I_{ZN} \in [0, 0.1]$$

From [2], the critical value is 1.96.

The proposed test for the example will be implemented as follows

Step-1: state the null hypothesis H_0 : there is no reduction in uncertainty of Covid-19 vs. the alternative hypothesis H_1 : there is a significant reduction in uncertainty of Covid-19.

Step-2: the value of the statistic is 2.42.

Step-3: Specify the level of significance $\alpha = 0.05$ and select the critical value from [2] which is 1.96.

Step-4: Do not accept the null hypothesis as the value of Z_N is larger than the critical value.

From the analysis, it can be seen that the calculated value of $Z_N \in [Z_L, Z_U]$ is larger than the critical value of 1.96. Therefore, the null hypothesis H_0 : there is no reduction in uncertainty of Covid-19 will be rejected in favor of H_1 : there is a significant reduction in uncertainty of Covid-19. Based on the study, it is concluded that there is a significant reduction in the uncertainty of Covid-19.

Simulation study

In this section, a simulation study is performed to see the effect of the measure of indeterminacy on statistic $Z_N \in [Z_L, Z_U]$. For this purpose, a neutrosophic form of $Z_N \in [Z_L, Z_U]$ obtained from the real data will be used. The neutrosophic form of $Z_N \in [Z_L, Z_U]$ is given as

$$Z_N = (1 + I_{ZN}) \frac{0.83 - 0.50}{\sqrt{\frac{0.50[1 - 0.50][1 - 0.50]}{(12 - 1)0.50}}}; I_{ZN} \in [I_{ZL}, I_{ZU}]$$

To analyze the effect on H_0 , the various values of $I_{ZN} \in [I_{ZL}, I_{ZU}]$ are considered. The computed values of $Z_N \in [Z_L, Z_U]$ along with the decision on H_0 are reported in Table 1. For this study $\alpha = 0.05$ and the critical value is 1.96. The null hypothesis H_0 will be accepted if the calculated value of Z_N is less than 1.96. From Table 1, it can be seen that as the values of $I_{ZN} \in [I_{ZL}, I_{ZU}]$ increases from 0.01 to 2, the values of $Z_N \in [Z_L, Z_U]$ increases. Although, a decision about H_0 remains the same at all values of measure of indeterminacy $I_{ZN} \in [I_{ZL}, I_{ZU}]$ but the difference between $Z_N \in [Z_L, Z_U]$ and the critical value of 1.96 increases as I_{ZU} increases. From the study, it can be concluded that the measure of indeterminacy $I_{ZN} \in [I_{ZL}, I_{ZU}]$ affects the values of $Z_N \in [Z_L, Z_U]$.

Table 1 The effect on $Z_N \in [Z_L, Z_U]$ for various values of $I_{ZN} \in [I_{ZL}, I_{ZU}]$

$I_{ZN} \in [I_{ZL}, I_{ZU}]$	$Z_N \in [Z_L, Z_U]$	Decision about H_0	$I_{ZN} \in [I_{ZL}, I_{ZU}]$	$Z_N \in [Z_L, Z_U]$	Decision about H_0
[0,0]	[2.20,2.20]	Reject H_0	[0,0.15]	[2.20,2.53]	Reject H_0
[0,0.01]	[2.20,2.22]	Reject H_0	[0,0.16]	[2.20,2.55]	Reject H_0
[0,0.02]	[2.20,2.24]	Reject H_0	[0,0.17]	[2.20,2.57]	Reject H_0
[0,0.03]	[2.20,2.26]	Reject H_0	[0,0.18]	[2.20,2.59]	Reject H_0
[0,0.04]	[2.20,2.28]	Reject H_0	[0,0.19]	[2.20,2.61]	Reject H_0
[0,0.05]	[2.20,2.31]	Reject H_0	[0,0.2]	[2.20,2.64]	Reject H_0
[0,0.06]	[2.20,2.33]	Reject H_0	[0,0.3]	[2.20,2.86]	Reject H_0
[0,0.07]	[2.20,2.35]	Reject H_0	[0,0.4]	[2.20,3.08]	Reject H_0
[0,0.08]	[2.20,2.37]	Reject H_0	[0,0.5]	[2.20,3.30]	Reject H_0
[0,0.09]	[2.20,2.39]	Reject H_0	[0,0.6]	[2.20,3.52]	Reject H_0
[0,0.10]	[2.20,2.42]	Reject H_0	[0,0.7]	[2.20,3.74]	Reject H_0
[0,0.11]	[2.20,2.44]	Reject H_0	[0,0.8]	[2.20,3.96]	Reject H_0
[0,0.12]	[2.20,2.46]	Reject H_0	[0,0.9]	[2.20,4.18]	Reject H_0
[0,0.13]	[2.20,2.48]	Reject H_0	[0,1]	[2.20,4.40]	Reject H_0
[0,0.14]	[2.20,2.50]	Reject H_0	[0,2]	[2.20,6.60]	Reject H_0

Comparative studies

As mentioned earlier, the proposed Z-test for uncertainty events is an extension of several tests. In this section, a comparative study is presented in terms of measure of indeterminacy, flexibility and information. We will compare the efficiency of the proposed Z-test for uncertainty with the proposed Z-test for uncertainty under classical statistics, proposed Z-test for uncertainty under fuzzy logic and proposed Z-test for uncertainty under interval statistics. The neutrosophic form of the proposed statistic $Z_N \in [Z_L, Z_U]$ is expressed as $Z_N = 2.20 + 2.20I_{ZN}; I_{ZN} \in [0, 0.1]$. Note that the first 2.20 presents the existing Z-test for uncertainty under classical statistics, the second part $2.20I_{ZN}$ is an indeterminate part and 0.1 is a measure of indeterminacy associated with the test. From the neutrosophic form, it can be seen that the proposed test is flexible as it gives the values of $Z_N \in [Z_L, Z_U]$ in an interval from 2.20 to 2.42 when $I_{ZU} = 0$. On the other hand, the existing test gives the value of 2.20. In addition, the proposed test uses information about the measure of indeterminacy that the existing test does not consider. Based on the information, the proposed test is interpreted as the probability that H_0 : there is no reduction in uncertainty of Covid-19 is accepted with a probability of 0.95, committing a type-I error is 0.05 with the measure of an indeterminacy 0.10. Based on the analysis, it is concluded that the proposed test is informative than the existing test. The proposed test is also better than the Z-test for uncertainty under fuzzy-logic as the test using fuzz-logic gives the value of the statistic from 2.20 to 2.42 without any information about the measure of indeterminacy. The test under interval statistic only considers the values within the interval rather than the crisp value. On the other hand, the analysis based on neutrosophic considers any type of set. Based on the analysis, it is concluded that the proposed Z-test is efficient than the existing tests in terms of information, flexibility, and indeterminacy.

Comparison using power of the test

In this section, the efficiency of the proposed test is compared with the existing test in terms of the power of the test. The power of the test is defined as the probability of rejecting H_0 when it is false and it is denoted by β . As mentioned earlier, the probability of rejecting H_0 when it is true is known as a type-I error is denoted by α . The values of $Z_N \in [Z_L, Z_U]$ are simulated using the classical standard normal distribution and neutrosophic standard normal distribution. During the simulation 100 values of $Z_N \in [Z_L, Z_U]$ are generated from a classical standard normal distribution and neutrosophic standard normal distribution with

mean $\mu_N = \mu_L + \mu_U I_{\mu_N}; I_{\mu_N} \in [I_{\mu_L}, I_{\mu_U}]$, where $\mu_L = 0$ presents the mean of classical standard normal distribution, $\mu_U I_{\mu_N}$ denote the indeterminate value and $I_{\mu_N} \in [I_{\mu_L}, I_{\mu_U}]$ is a measure of indeterminacy. Note that when $I_{\mu_L} = 0$, μ_N reduces to μ_L . The values of $Z_N \in [Z_L, Z_U]$ are compared with the tabulated value at $\alpha = 0.05$. The values of the power of the test for the existing test and for the proposed test for various values of I_{μ_U} are shown in Table 2. From Table 2, it is clear that the existing test under classical statistics provides smaller values of the power of the test as compared to the proposed test at all values of I_{μ_U} . For example, when $I_{\mu_U} = 0.1$, the power of the test provided by the Z-test for uncertainty events under classical statistics is 0.94 and the power of the test provided by the proposed Z-test for uncertainty events is 0.96. The values of the power of the test for Z-test for uncertainty events under classical statistics and Z-test for uncertainty events under neutrosophic statistics are plotted in Fig. 1. From Fig. 1, it is quite clear that the power curve of the proposed test is higher than the power curve of the existing test. Based on the analysis, it can be concluded that the proposed Z-test for uncertainty events under neutrosophic statistics is efficient than the existing Z-test for uncertainty events.

Conclusions

The Z-test of uncertainty was introduced under neutrosophic statistics in this paper. The proposed test was a generalization of the existing Z-test of uncertain events under classical statistics, fuzzy-based test, and interval statistics. The performance of the proposed test was compared with the listed existing tests. From the real data and simulation study, the proposed test was found to be more efficient in terms of information and power of the test. Based on the information, it is recommended to apply the proposed test to check the reduction in

Table 2 The values of the power of two tests

I_{μ_U}	Existing Test	The Proposed Test
	$1 - \beta$	$1 - \beta$
0.05	0.96	0.97
0.1	0.94	0.96
0.2	0.94	0.96
0.3	0.91	0.97
0.4	0.93	0.95
0.5	0.89	0.95
0.75	0.89	0.92
1	0.93	0.96




Fig. 1 The power curves of the two tests

uncertainty under an indeterminate environment. The proposed test for big data can be considered as future research. The proposed test using double sampling can also be studied as future research. The estimation of sample size and other properties of the proposed test can be studied in future research.

Acknowledgements

We are thankful to the editor and reviewers for their valuable suggestions to improve the quality of the paper.

Author's contributions

MA wrote the paper.

Funding

None

Availability of data and materials

All data generated or analysed during this study are included in this published article

Declarations

Ethics approval and consent to participate

N/A

Consent for publication

N/A

Competing interests

None

Received: 27 September 2021 Accepted: 31 March 2022

Published online: 06 April 2022

References

1. DOLL H, CARNEY S. Statistical approaches to uncertainty: p values and confidence intervals unpacked. *BMJ evidence-based medicine*. 2005;10(5):133–4.
2. Kanji, G.K. *100 statistical tests*2006: Sage.
3. Lele SR. How should we quantify uncertainty in statistical inference? *Front Ecol Evol*. 2020;8:35.
4. Wang F, et al. Re-evaluation of the power of the mann-kendall test for detecting monotonic trends in hydrometeorological time series. *Front Earth Sci*. 2020;8:14.
5. Maghsoodloo S, Huang C-Y. Comparing the overlapping of two independent confidence intervals with a single confidence interval for two normal population parameters. *J Stat Plan Inference*. 2010;140(11):3295–305.
6. Rono BK, et al. Application of paired student t-test on impact of anti-retroviral therapy on CD4 cell count among HIV Seroconverters in serodiscordant heterosexual relationships: a case study of Nyanza region. Kenya.
7. Zhou X-H. Inferences about population means of health care costs. *Stat Methods Med Res*. 2002;11(4):327–39.
8. Niwitpong S, Niwitpong S-a. Confidence interval for the difference of two normal population means with a known ratio of variances. *Appl Math Sci*. 2010;4(8):347–59.

9. Viertl R. Univariate statistical analysis with fuzzy data. *Comput Stat Data Anal.* 2006;51(1):133–47.
10. Filzmoser P, Viertl R. Testing hypotheses with fuzzy data: the fuzzy p-value. *Metrika.* 2004;59(1):21–9.
11. Tsai C-C, Chen C-C. Tests of quality characteristics of two populations using paired fuzzy sample differences. *Int J Adv Manuf Technol.* 2006;27(5):574–9.
12. Taheri SM, Arefi M. Testing fuzzy hypotheses based on fuzzy test statistic. *Soft Comput.* 2009;13(6):617–25.
13. Jamkhaneh EB, Ghara AN. Testing statistical hypotheses with fuzzy data. In: 2010 International Conference on Intelligent Computing and Cognitive Informatics: IEEE; 2010.
14. Chachi J, Taheri SM, Viertl R. Testing statistical hypotheses based on fuzzy confidence intervals. *Austrian J Stat.* 2012;41(4):267–86.
15. Kalpanapriya D, Pandian P. Statistical hypotheses testing with imprecise data. *Appl Math Sci.* 2012;6(106):5285–92.
16. Parthiban, S. and P. Gajivaradhan, A Comparative Study of Two-Sample t-Test Under Fuzzy Environments Using Trapezoidal Fuzzy Numbers.
17. Montenegro M, et al. Two-sample hypothesis tests of means of a fuzzy random variable. *Inf Sci.* 2001;133(1-2):89–100.
18. Park S, Lee S-J, Jun S. Patent big data analysis using fuzzy learning. *Int J Fuzzy Syst.* 2017;19(4):1158–67.
19. Garg H, Arora R. Generalized MacLaurin symmetric mean aggregation operators based on Archimedean t-norm of the intuitionistic fuzzy soft set information. *Artif Intell Rev.* 2020:1–41.
20. Smarandache F. Neutrosophy. Neutrosophic probability, set, and logic, *ProQuest Information & Learning*, vol. 105. Michigan: Ann Arbor; 1998. p. 118–23.
21. Broumi S, Smarandache F. Correlation coefficient of interval neutrosophic set. In: *Applied mechanics and materials: Trans Tech Publ*; 2013.
22. Abdel-Basset M, et al. A novel group decision making model based on neutrosophic sets for heart disease diagnosis. *Multimed Tools Appl.* 2019:1–26.
23. Alhasan KFH, Smarandache F. Neutrosophic Weibull distribution and Neutrosophic Family Weibull Distribution 2019. Infinite Study.
24. Das SK, Edalatpanah S. A new ranking function of triangular neutrosophic number and its application in integer programming. *Int J Neutrosophic Sci.* 2020;4(2).
25. El Barbary G, O. and R. Abu Gdairi, Neutrosophic logic-based document summarization. *J Undergrad Math.* 2021.
26. Smarandache, F., *Introduction to neutrosophic statistic* 014: Infinite Study.
27. Chen J, Ye J, Du S. Scale effect and anisotropy analyzed for neutrosophic numbers of rock joint roughness coefficient based on neutrosophic statistics. *Symmetry.* 2017;9(10):208.
28. Chen J, et al. Expressions of rock joint roughness coefficient using neutrosophic interval statistical numbers. *Symmetry.* 2017;9(7):123.
29. Sherwani RAK, et al. A new neutrosophic sign test: an application to COVID-19 data. *PLoS One.* 2021;16(8):e0255671.
30. Aslam M. Neutrosophic statistical test for counts in climatology. *Sci Rep.* 2021;11(1):1–5.
31. Albassam M, Khan N, Aslam M. Neutrosophic D'Agostino test of normality: an application to water data. *J Undergrad Math.* 2021;2021.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

